

**Table 55. Five-year relative cancer survival rates for selected cancer sites, according to race and sex:
Selected geographic areas, 1974–79, 1980–82, 1983–85, 1986–88, 1989–91, and 1992–98**

[Data are based on the Surveillance, Epidemiology, and End Results Program's 9 population-based cancer registries]

Sex and site	White						Black or African American					
	1974–79	1980–82	1983–85	1986–88	1989–91	1992–98	1974–79	1980–82	1983–85	1986–88	1989–91	1992–98
Both sexes												Percent of patients
All sites	50.9	52.1	53.9	56.7	60.3	63.8	39.2	39.7	39.8	42.6	46.2	52.6
Oral cavity and pharynx	54.9	55.6	55.3	55.2	55.4	58.8	36.5	31.0	35.2	34.7	32.6	34.9
Esophagus	5.5	7.3	9.3	10.7	11.8	14.7	3.3	5.4	6.2	7.2	8.9	8.3
Stomach	15.2	16.5	16.2	19.1	18.4	20.9	15.7	19.4	18.9	19.3	24.8	20.0
Colon	51.9	55.7	58.4	61.6	63.1	62.7	47.3	49.3	49.3	52.9	53.9	52.8
Rectum	49.8	53.1	55.9	59.1	60.5	62.4	40.4	37.9	43.9	51.1	54.3	52.7
Pancreas	2.4	2.8	2.9	3.1	4.1	4.3	3.2	4.5	5.0	6.2	3.8	3.9
Lung and bronchus	13.1	13.5	13.8	13.5	14.3	15.0	11.3	12.1	11.3	11.9	10.7	12.3
Urinary bladder	74.9	78.9	78.2	80.7	82.1	82.3	51.5	58.3	59.6	62.3	61.9	64.5
Non-Hodgkin's lymphoma	48.2	51.9	54.4	52.9	51.9	56.1	50.3	50.2	45.1	50.2	43.7	46.1
Leukemia	36.6	39.5	42.0	44.2	45.8	47.3	30.8	32.9	33.6	38.0	34.1	38.4
Male												
All sites	43.4	46.7	48.5	51.8	57.7	63.5	32.1	34.3	34.5	37.7	43.3	54.0
Oral cavity and pharynx	54.4	54.6	54.4	52.2	52.1	57.8	31.2	26.5	29.9	29.3	27.9	29.5
Esophagus	5.2	6.5	7.8	11.2	11.8	14.7	2.3	4.6	5.0	7.0	8.0	9.0
Stomach	13.8	15.6	14.6	16.2	15.1	19.5	15.3	18.5	18.5	15.3	22.3	19.2
Colon	51.0	56.0	58.9	62.5	63.7	63.0	45.4	46.7	48.3	52.6	53.8	53.2
Rectum	48.9	51.7	55.2	58.8	60.2	61.6	36.7	36.1	43.1	46.2	56.0	51.6
Pancreas	2.6	2.5	2.7	2.9	3.8	4.3	2.4	3.2	4.4	6.4	3.3	4.2
Lung and bronchus	11.6	12.2	12.1	12.0	12.7	13.3	10.0	10.9	10.3	11.9	9.5	10.8
Prostate gland	70.3	74.5	76.3	82.7	91.8	97.8	60.8	64.7	63.9	69.3	80.6	92.6
Urinary bladder	75.9	79.9	79.5	82.3	84.4	84.4	58.9	62.8	64.8	67.4	65.5	69.4
Non-Hodgkin's lymphoma	47.2	50.9	53.4	50.1	47.7	52.7	44.7	47.2	43.6	46.6	38.7	41.4
Leukemia	35.6	39.6	41.6	45.5	46.4	48.4	30.7	30.1	32.5	36.7	29.2	37.9
Female												
All sites	57.4	57.1	58.8	61.5	62.9	64.2	46.9	46.0	45.4	47.9	49.3	51.0
Colon	52.6	55.4	58.0	60.7	62.5	62.4	48.7	51.3	50.1	53.1	53.9	52.5
Rectum	50.7	54.7	56.7	59.5	60.8	63.5	43.8	40.2	44.7	56.0	52.6	53.8
Pancreas	2.3	3.0	3.2	3.4	4.4	4.4	4.1	5.8	5.5	6.0	4.3	3.7
Lung and bronchus	16.7	16.2	17.0	15.8	16.6	17.0	15.5	15.5	14.1	11.7	13.0	14.7
Melanoma of skin	85.8	88.3	89.4	91.3	91.5	91.8	69.9	*	70.1	*	94.0	61.9
Breast	75.4	77.1	79.2	83.9	86.1	87.6	63.1	65.7	63.4	69.2	71.1	72.5
Cervix uteri	69.7	68.2	70.6	71.7	72.3	72.1	63.0	61.3	60.4	55.5	62.6	59.9
Corpus uteri	87.7	82.8	84.5	84.4	85.6	86.0	59.4	55.1	54.2	57.1	57.5	60.5
Ovary	37.2	38.7	40.2	41.9	49.6	52.5	40.5	39.1	41.6	38.6	41.6	52.5
Non-Hodgkin's lymphoma	49.3	52.9	55.5	56.2	57.1	60.2	57.5	53.7	46.9	54.8	50.8	53.8

* Data for population groups with fewer than 25 cases are not shown.

NOTES: Rates are based on followup of patients through 1999. The rate is the ratio of the observed survival rate for the patient group to the expected survival rate for persons in the general population similar to the patient group with respect to age, sex, race, and calendar year of observation. It estimates the chance of surviving the effects of cancer. The race groups white and black include persons of Hispanic and non-Hispanic origin. Numbers have been revised and differ from previous editions of *Health, United States*.

SOURCE: National Institutes of Health, National Cancer Institute, Surveillance, Epidemiology, and End Results (SEER) Program at www.seer.cancer.gov.